

Towards the Final State of Spherical Gravitational Collapse and Likely Source of Gamma Ray Bursts

Abhas Mitra

Theoretical Physics Division, Bhabha Atomic Research Center,
Mumbai-400085, India

E-mail: amitra@apsara.barc.ernet.in

By analyzing the *global properties* of the equations for general relativistic collapse which reflect the rate of change of the *circumference radius* R with the *proper radius* l , we show that (i) no “trapped surface” is ever formed and (ii) the *gravitational mass* of the fluid $M(r, t) \rightarrow 0$ as $R \rightarrow 0$. This result is absolutely independent on the equation of state of the fluid or any other details. Thus, for continued collapse, the total energy output measured by a distant observer, is $Q \rightarrow M_i c^2$, the original gravitational mass.

PACS numbers: 04.20.Dw, 04.70. Bw

The problem of gravitational collapse has been described as “the greatest crisis in physics of all time” [1](pp.1196). Though the theoretical framework for studying this problem is fairly well developed [2,3], in practice it is impossible to make much headway without making a number of simplifying assumptions because of our inability to *self consistently* handle: (i) the equation of state (EOS) of matter at arbitrarily high density and temperature, (ii) the opacity of nuclear matter at such likely unknown extreme conditions, (iii) the associated radiation transfer problem and all other highly nonlinear and coupled partial differential equations. As far as exact solution is concerned, it is possible only if the collapsing fluid is considered to be a *dust* with pressure $p \equiv 0$ everywhere including the center even for *arbitrary high density*. Further, it should be considered *homogeneous*, and the pioneering solution to this effect was obtained by Oppenheimer and Snyder [4](OS) which suggested that the spherical dust ball collapses to a point to a state of infinite proper density, i.e., $R \rightarrow 0$, $\rho \rightarrow \infty$ in a *proper time* (*by fiat, pressure still remains zero*):

$$\tau = \frac{\pi}{2} \left(\frac{3}{8\pi\rho(0)} \right)^{1/2} \quad (1)$$

where $\rho(0)$ is the initial mass-energy density of the fluid when it is assumed to be *at rest*. A distant inertial observer S_∞ , however can not witness this approach to a physical singularity because as $R \rightarrow R_{gi} = 2GM_i/c^2$, where M_i is the initial gravitational mass of the dust, the photons emitted from the surface of the collapsing fluid are infinitely redshifted indicating the formation of an “event horizon” and a subsequent black hole (BH). Attempts for numerical studies [5] to study the collapse of a physical fluid can progress only by presuming that

the energy emitted during the process $Q \ll M_i c^2$. The said assumption tantamounts to the condition that the eventual temperature of the nuclear matter $T \ll 1\text{GeV}$, the typical nucleon mass. When this is so, one can use a $T = 0$ degenerate EOS or can incorporate a finite T EOS which incorporates a first order correction term $\propto T^2$. A general overall self-consistent numerical treatment of the problem which does not impose explicit or tacit smallness conditions on Q and T is implausible. For a dust, thermodynamics would tell that a $p = 0$ condition implies that the internal energy density $e = 0$ and also $T = 0$. Consequently, the instantaneous value of $Q \equiv 0$, which in turn implies the instantaneous gravitational mass of the body $M \equiv M_i$. The presumption that $Q \ll M_i c^2$ leads to a more or less similar condition, that $M \approx M_i$, and naturally numerical computations reveal the formation of an event horizon enclosing a mass $M_g = M_f \approx M_i$ within a finite proper time. However, when the condition that for all times $Q \ll M_i c^2$ is not imposed the running value of $M = M_i - Q/c^2 < M_i$, consequently, the would be value of R_g will be smaller than the initially suspected value of R_{gi} leading to value of Q higher than the previous value. This may in turn lead to a value of M lower than its previously expected value. And, in general, it can not be predicted whether this apparently runaway process converges to a certain finite value of M_g in a finite time or not, where M_g is the mass at $R = R_g$. In this situation, we would like to explore the limiting properties of the equations for collapse for a perfect fluid *without making any conceptual assumption*. The matter part of the fluid has the energy momentum tensor

$$T^{\mu\nu} = (p + \rho)u^\mu u^\nu + pg^{\mu\nu} \quad (2)$$

where u^μ is the fluid 4-velocity, and $\rho = \rho_0 + e$, with $\rho_0 = mn$, m is the rest mass of a baryon as measured by S_∞ and n is the proper number density of the baryons. Now we have taken $c = 1$. The radiation part of the fluid is described by

$$E^{\mu\nu} = qk^\mu k^\nu \quad (3)$$

where q is both the energy density and the radiation flux in the proper frame and $k^\mu = (1; 1, 0, 0)$ is a null geodesic vector [6] so that $k^\mu k_\mu = 0$. Irrespective of whether a “trapped surface” is formed or not, the interior of the collapsing body may be described by the Schwarzschild metric [1–3,6]:

$$ds^2 = -e^{2\phi} dt^2 + e^{2\lambda} dr^2 + R(r, t)^2 (d\vartheta^2 + \sin^2 \vartheta d\varphi^2) \quad (4)$$

Here r is a comoving coordinate, so that the total number of baryons $N(r)$ enclosed within the sphere of constant coordinate radius r is constant and independent of t and $R(r, t)$ is the circumference coordinate. The explicit forms of e^ϕ and e^λ are not known, but they are nonsingular for any finite R , and computational techniques are supposed to self-consistently evaluate them. The exterior region must be described in terms of Vaidya's retarded time u [7], which turns out to be the proper time of S_∞ [3]:

$$ds^2 = -e^{2\psi} du^2 - 2e^\psi e^\lambda du dr + R^2 (d\vartheta^2 + \sin^2 \vartheta d\varphi^2) \quad (5)$$

where $e^{2\psi} = (1 - 2GM(u)/R)$. The comoving time and radial derivative operations are respectively defined by

$$D_t = e^{-\phi} \left(\frac{\partial}{\partial t} \right)_r = \frac{d}{d\tau}; \quad D_r = e^{-\lambda} \left(\frac{\partial}{\partial r} \right)_t = \frac{d}{dl} \quad (6)$$

Here $d\tau = e^\phi dt$ is an element of proper time at constant r , and $dl = e^\lambda dr$ is an element of proper radial distance between two baryon-shells r and $r+dr$, ϑ and φ are frozen in spherical symmetry. The *circumference velocity* of the fluid measured at constant r in terms of proper time [2,3] is

$$D_t R = U = \frac{dR}{d\tau} \quad (7)$$

Here D_t and D_r are to be treated as equivalents of "convective derivative" following the trajectory of the fluid parcel used in classical hydrodynamics. The central *global character* of the collapse process, *valid in both metrics*, is manifest through

$$\Gamma = D_r R = \frac{d(\text{circumference})/2\pi}{d(\text{proper distance})} = \frac{dR}{dl} \quad (8)$$

And the most important *global constraint* on the field equations is that the relationship

$$\Gamma^2 = 1 - \frac{2GM}{R} + U^2 \quad (9)$$

is *preserved for all times* [2,3]. The actual equations for containing the details of the collapse process, and *valid in metric* (4), are [3]

$$D_r M = 4\pi R^2 [\Gamma(\rho + q) + Uq] \quad (10)$$

$$D_t M = -4\pi R^2 pU - L(U + \Gamma) \quad (11)$$

$$D_t U = -\frac{\Gamma}{\rho + p} \left(\frac{\partial p}{\partial R} \right)_t - \frac{M + 4\pi R^3 (p + q)}{R^2} \quad (12)$$

$$D_t \Gamma = -\frac{U\Gamma}{\rho + p} \left(\frac{\partial p}{\partial R} \right)_t + \frac{L}{R} \quad (13)$$

where the comoving luminosity is

$$L = 4\pi R^2 q \quad (14)$$

For motionless matter in flat space (S_∞), $\Gamma \equiv 1$. In a curved space $\Gamma < 1$ because of the gravitational field. And a value of $\Gamma < 0$ would signify that the spacetime is pinched off from the rest of the universe [3], and this is what is believed to happen in the final stages of gravitational collapse. However *nobody has ever hinted what could be order of magnitude of the (-ve) value of final Γ_f even for the idealized case of a dust collapse*. The popular conviction that even a physical fluid must eventually behave like a O-S dust is based on certain properties of the *radiationless* collapse equations [2]:

$$D_r M = 4\pi R^2 \rho \Gamma \quad (15)$$

$$D_t M = -4\pi R^2 pU \quad (16)$$

$$D_t U = -\frac{\Gamma}{\rho + p} \left(\frac{\partial p}{\partial R} \right)_t - \frac{M + 4\pi R^3 p}{R^2} \quad (17)$$

$$D_t \Gamma = -\frac{U\Gamma}{\rho + p} \left(\frac{\partial p}{\partial R} \right)_t \quad (18)$$

The traditional argument favouring BH formation goes like this [8]: For collapse, $U < 0$, and from eq. (16) it follows that $M(r, t)$ increases monotonically within the fluid; of course, it remains constant, for adiabatic collapse, for $R \geq R_0$, the outer boundary. Because of this, the M/R^2 term in eq. (17) increases more rapidly than it would in the Newtonian theory where it remains constant. Then eq.(18) would suggest that Γ decreases monotonically instead of remaining constant in a Newtonian case. Both of these relativistic effects are supposed to cause the collapse process monotonic and accelerate faster than the Newtonian case [8]. Here note that unless we presume that $\partial p/\partial R$ can be positive, a negative value of Γ in eq.(18) would demand that Γ increases with time implying Γ would tend to be back to zero. In any case when we restore q , or in other words, when we do away with the assumption $Q \ll M_i$, *this entire line of argument collapses* and nothing can be predicted about the nature of evolution of the collapse process. In this situation, we discuss below the global properties of these equations. Following the discussion by Landau & Lifshitz [9](pp.250), the *ordinary radial velocity of the fluid as measured in a local rest frame* is

$$v^{\hat{r}} \equiv v = \frac{e^\lambda dr}{e^\phi dt} \equiv \frac{dl}{d\tau} \quad (19)$$

By **principle of equivalence** (POE), $v \leq c$. By combining eq. (7), (8), and (20), we find

$$U = \Gamma v \quad (20)$$

The above relationship does not depend on the signature or form of the metric coefficients and, as mentioned earlier, all equations involving *interior Sch. metric coefficients* remain valid even if a trapped surface is formed [6]. Further, eq. (20) *must not be confused with* $u^{\hat{r}} = \gamma v$, where $u^{\hat{r}}$ is the radial component of the 4-velocity measured in the local inertial frame (in the comoving Lagrangian frame, $u^r \equiv 0$) and $\gamma = (1 - v^2)^{-1/2}$ is the Lorentz Factor of the fluid in the same inertial frame. By combining eq. (9) and (19), we obtain

$$\frac{\Gamma^2}{\gamma^2} \equiv 1 - \frac{2GM}{R} \quad (21)$$

Since the LHS of the foregoing eq. is positive (even if Γ were negative), so will be the RHS, and therefore, **by simply using POE**, we find that

$$\frac{2GM}{R} \leq 1 \quad (22)$$

Recall that, for spherical collapse the condition for formation of a *trapped surface*, a surface of no-return is $2GM/R > 1$ [3,10]. Thus, in a most general manner, we find that *trapped surfaces are not allowed by GTR* contrary to the assumption made in a large number of previous works starting from the sixties. It is the assumption (1) about the occurrence of the trapped surfaces, along-with, (2) the causality conditions requiring the nonexistence of closed timelike lines, (3) the strong energy condition ($T_{\mu\nu} - 0.5g_{\mu\nu}u^\mu u^\nu \geq 0$, and (4) a certain generality condition on the Riemann-Christoffel tensor, that laid the foundation of the famous singularity theorems of Hawking, Penrose, Geroch and others [10]. So now they are invalidated because trapped surfaces are not allowed by GTR. We quickly recall the relevant comments by Misner et al. [1](pp.935): “All the conditions except the trapped surface seem eminently reasonable for any physically realistic space time”. Thus, the mathematical and moral foundation for the inevitability of occurrence of singularities in GTR are eliminated. However, this does not necessarily mean that singularities must not occur, and, we will probe this question in an accompanying paper [10]. Any EOS will support only a certain finite maximum M_i and therefore given sufficiently large M_i , continued collapse is a genuine possibility; and if indeed $R \rightarrow 0$, then in order to satisfy the foregoing inequality, *we must have*

$$M_f \rightarrow 0; \quad R_f \rightarrow 0 \quad (23)$$

Remember here that the quantity $M_0 = mN$ (which is the baryonic mass of the star, if there are no antibaryons)

is conserved as $M_f \rightarrow 0$. How do reconcile our general result $M_f = 0$, with the result of O-S collapse which suggests that as the dust ball with an initial condition $R = R_0$, $M(r_0, 0) = M_i$ collapses from a *state of rest*, i.e., $v = v(r, 0) = 0$ and $U = U(r, 0) = 0$, it forms a BH with $M_f = M_i$ in a finite proper time? To find the $\rho(0)$ occurring in eq. (1) in a truly self-consistent way, we must fix it from the collapse equations themselves, rather than *artificially and by hand*. When we put this $U = q = 0$ condition in eq.(12), we obtain the so-called Oppenheimer-Volkov (OV) equation [11].

$$\frac{dp}{dR} = -\frac{p + \rho(0)}{R(R - 2GM)}(4\pi p R^3 + 2GM) \quad (24)$$

By feeding the $p = 0$ EOS in the OV equation it may be verified that whether the fluid is considered homogeneous or not the solutions could be of two types: (i) $\rho(0) = 0$, $R_0 = \infty$, $M_i = \text{finite}$ and (ii) $\rho(0) = 0$, $R_0 = \text{finite}$, $M_i = 0$. In the former case, the fluid boundary requires infinite τ for attaining a finite R or R_g , implying that the collapse process hardly ever takes off:

$$\tau(R) = \frac{2}{3} \left[\frac{R_0}{c} \left(\frac{R_0}{R_g} \right)^{1/2} - \frac{R}{c} \left(\frac{R}{R_g} \right)^{1/2} \right] \quad (25)$$

and there is no question of formation of a BH. And in the latter case, our proof that $M_f \equiv 0$ is vindicated because $M_f = M_i = 0$. Even in this case, eq. (1) yields $\tau = \infty$! These **boundary conditions** are *fudged in all theoretical and numerical discussions by pretending that a self-supporting fluid with finite values of ρ and R_0 , and $p \neq 0$, everywhere, may approximately be considered to suddenly attain the dust-like dynamics*. Even if such a dust-like collapse took place, it can be verified from eq.(18) (by evolving from $U = 0$ condition) that, far from assuming any negative value, the value of Γ remains fixed to the initial value Γ_0 , which is unity, and which is the appropriate value of Γ for a motionless (no kinetic energy) and matter free (thus gravityfree) region. It might be thought that if the numerical calculations study the collapse of a supermassive star, the horizon may be formed at very small densities $\sim 10^{-4} \text{g/cm}^3$, and the entire collapse physics including the EOS can be handled accurately. This idea would not be correct because, the expectation that horizon would be formed at $R_g \approx 2GM_i$ is based on the presumption that $Q \ll M_i$. And in any case our central result manifest by eq. (22) is an exact one. In the context of a numerical neutron star collapse, we will discuss about an ansatz for handling the thermodynamics of the problem which may reveal a value of Q much higher than previously reported values, although the eventual computation must break down as T_{com} keeps on increasing without any upper limit [13].

It is widely believed that Chandrasekhar’s discovery that White Dwarfs (WD) can have a maximum mass set

the stage for having a gravitational singular state with finite mass. The hydrostatic equilibrium of WDs can be approximately described by Newtonian polytropes [14] for which one has $R \propto \rho_c^{(1-n)/2n}$, where ρ_c is the central density of the polytrope having an index n . It shows that, for a singular state i.e., for $\rho_c \rightarrow \infty$, one must have $R \rightarrow 0$ for $n > 1$; and Chandrasekhar's limiting WD indeed has a *zero radius*. On the other hand, the mass of the configuration $M \propto \rho_c^{(3-n)/2n}$. And unless $n = 3$, $M \rightarrow \infty$ for the singular state. However, for ultrarelativistic electrons, the EOS is $p \rightarrow e/3$ and the corresponding $n \rightarrow 3$ and one obtains a finite value of M_{ch} - the Chandrasekhar mass. Now when we apply the theory of polytropes for a case where the pressure is supplied by the baryons and not only by electrons, we must consider GTR polytropes [15]. It can be easily verified from eq. (2.24) of ref.(15) that in the limit $\rho_c \rightarrow \infty$, the scale size of GTR polytropes $A^{-1} \rightarrow 0$. Further eq.(2.15) and (4.7) of ref. (15) tell that $M \propto K^{n/2} \propto \rho_c^{-1/2} \rightarrow 0$ for $\rho_c \rightarrow \infty$. Thus, a *proper GTR extension of Chandrasekhar's work would not lead to a BH of finite mass*, but, on the other hand, to a singular state with $M \rightarrow 0$. Physically, the $M = 0$ state may result when the *negative gravitational energy* exactly cancels the internal energy, the *baryonic mass energy* M_0 and any other energy, and which is possible in the limit $\rho \rightarrow \infty$ and $p \rightarrow \infty$. In fact, Zeldovich and Novikov [12] discussed the possibility of having an ultracompact configuration of degenerate fermions obeying the EOS $p = e/3$ with $M \rightarrow 0$; we have verified that their configuration actually corresponds to $R \rightarrow R_g \rightarrow 0$. In fact, they mentioned the possibility of *having a machine for which $Q \rightarrow M_i c^2$* . In a different context, it has been argued that *naked singularities* produced in spherical collapse must have $M_f = 0$ [16]. Also, it was discussed long ago [15] that spherical gravitational collapse should come to a *decisive end* with $M_f = M^* = 0$, and, in fact, this understanding was formulated as a "Theorem" [17]:

"THEOREM 23. Provided that matter does not undergo collapse at the microscopic level at any stage of compression, then, -regardless of all features of the equation of state - there exists for each fixed number of baryons A a "gravitationally collapsed configuration", in which the mass-energy M^* as sensed externally is zero." We have proved this and more for spherical symmetry, our result is absolutely general and has got nothing to with the exact EOS and radiation transport equations. On the other hand, it depends only on the global properties of the GTR collapse equations and the Principle of Equivalence. We have learnt that GRB990123 released an energy 3×10^{54} ergs of energy in γ -rays alone under condition of isotropy. Given the fact that the relativistic blast wave radiating the gamma-rays may not have an efficiency greater than 10-20% in converting the blast wave energy into gamma rays, the total electromagnetic energy released in the process could easily be $Q_{FB} \sim 10^{55}$

erg. Further, the energy output in ν s could be or rather should be $> Q_{FB}$, the net energy released in the process could be $Q \sim \text{few } 10^{55}$ erg. In case, there is weak beaming, the total energy released may still be $Q \leq 10^{55}$ erg. physical mechanism, other than what has been proposed here, can satisfactorily explain the origin of such unprecedented energy. Such energy release is hardly possible if trapped surfaces would form.

Acknowledgement: I am thankful to Prof. P.C. Vaidya for painstakingly going through this work, offering some suggestions, and confirming that this work is physically and general relativistically correct.

-
- [1] C.W. Misner, K.S. Thorne, and J.A. Wheeler, *Gravitation* (W.H. Freeman and Company, San Fransisco, 1973)
 - [2] For adiabatic collapse eq. see C.W. Misner and D.H. Sharp, Phys. Rev. **136** **2B**, 571 (1964); also M. May & R. White, Phy. Rev. **141**, 1233 (1966).
 - [3] For physical collapse eq. see C.W. Misner, Phys. Rev. **137** **B**, 1360 (1965); R.W. Lindquist, R.A. Schwartz, & C.W. Misner, Phy. Rev. **137** **B**, 1364 (1965); W.C. Hernandez & C.W. Misner, Astrophys. J. **143**, 452 (1966), C.W. Misner, in Relativity Theory and Astrophysics (Vo. 3), ed. J.E. Ehlers, see p.117-128, and P.C. Vaidya, Astrophys. J., **144**, 943 (1965).
 - [4] J.R. Oppenheimer & H. Snyder, Phy. Rev., **56**, 455 (1939); however, the collapse of an inhomogeneous dust was already discussed by B. Datt, Phys. Zeits. bf 108, 341 (1938). In general the inhomogeneous dust case may not be amenable to analytical treatment and there is a raging debate whether one can have a "naked singularity" in such cases.
 - [5] T.W. Baumgarte et al., Astrophys. J., **468**, 823 (1996), T. Baumgarte, S.L. Shapiro, & S.A. Tekuolsky, Astrophys. J., **458**, 680 (1996), W. Keil & H.T. Janka, Astron. Astrophys., **296**, 145 (1995). E. Gourgoulhon & P. Haensel, Astron. Astrophys., **271**, 209 (1993). It may be noted that none of the recent and *more accurate computations report the occurrence of a negative Γ region*.
 - [6] P.C. Vaidya, (1998), Personal communication.
 - [7] P.C. Vaidya, Nature, **171**, 260 (1953).
 - [8] J.C. Miller and D.W. Sciama, in General Relativity and Gravitation, Vol. 2., ed. A. Held (Plenum, New York, 1980). see p. 369.
 - [9] L.D. Landau & E.M. Lifshitz, *The Classical Theory of Fields*, (Pergamon Press, Oxford, 1975, 4th ed.), see p. 286, 311, 316.
 - [10] S.W. Hawking and G.R.F. Ellis, *The Large Scale Structure of the Universe* (Cambridge Univ. Press, Cambridge, 1973), pp. 261-275.
 - [11] J.R. Oppenheimer & G.M. Volkov, Phy. Rev. **55**, 375 (1939).
 - [12] Ya. B. Zeldovich & I.D. Novikov, *Relativistic Astrophysics*, Vol. 1 (Univ. Chicago Press, Chicago, 1971), see

pp. 95 and pp. 297.

- [13] Present author, in preparation.
- [14] S. Chandrasekhar, *An Introduction To The Study of Stellar Structure*, (Dover, New York, 1967), see eq. 62, 69, p.97.
- [15] R.F. Tooper, *Astrophys. J.* **140**, 393 (1964).
- [16] K. Lake, *Phys. Rev. Lett.* **68**, 3129 (1992)
- [17] B.K. Harrison, K.S. Thorne, M. Wakano, & J.A. Wheeler, *Gravitation Theory & Gravitational Collapse*, (Univ. Chicago, Chicago, 1965), see p. 75.